



Calf Xtreme instructions for use

- Mix with cold or lukewarm water or milk in the protein shake provided - **Do not use hot water!!!**
- Stir thoroughly with milk and feed out within 2-3 hours. (milk will start to curdle if stored too long - especially if added to warm milk)
- **Dose Rate**
- Standard dose rate: 1gm/calf/day
- Stress events (Bad weather, change of diet and debudding). Double the dose for a couple of days either side: 2gm/calf/day.
- For most disease challenges including Rotavirus. 2gm/calf/day in milk once a day, with electrolytes fed in a second feed.
- For Cryptosporidium and Salmonella. 4gm/calf/day in milk once a day with electrolytes fed in a second feed.
- Calf Xtreme Plus can be used in conjunction with antibiotics so the disease challenge should be diagnosed and appropriate antibiotics also used.
- For scouring calves a binding agent should also be used to slow the passage of dung in the lower gut so that the bacteria in Calf Xtreme are able to work.
- Do not use a bacterial desiccant with Calf Xtreme
- If feeding low quantities of milk or milk powder (under 3L), split Calf Xtreme dose between feeds

FOR ANY ANIMAL HEALTH CONCERNS PLEASE CONSULT A VET.

FOR ANY QUESTIONS REGARDING CALF XTREME, CONTACT US

CHRIS - 027 4591 061

MATT - 021 234 1713

Calf Xtreme suggested feed rates



Note: These are feeding guidelines based on what is possible and beneficial using Calf Xtreme. However observations must still be made and do not exceed what your calves can comfortably intake

Ingredients -

P7 Active dry yeast
5 strains of probiotic bacteria
5 digestive enzymes

Whole milk:

Friesian

- 4L TAD for 3 - 7 days then 6-7L OAD thereafter.

Kiwicross

- 3L TAD for 3 - 7 days then 5-6L OAD thereafter.

Jersey

- 2-2.5L TAD for 7-10 days then 4-5 L OAD thereafter

Wagyu

- 2-2.5L TAD for 7-10 days then 4-5 L OAD thereafter

Calf milk replacer:

Follow manufacturers instructions for the first 2-3 weeks then feed once a day increasing volume offered to the calf using the twice a day feeding instructions e.g. Where instructions for twice a day suggest 150g/L, feed 750g/5L or 900g/6L over 1 feed. Alternatively, contact us and we can talk this through.

Note - While we are not aware of the ability to overdose on Calf Xtreme, we don't recommend using more than 10g in one feed or multiplying the dose for more than 5 days. If feeding low volumes of 3L or less, split the dose between 2 feeds!

FOR ANY ANIMAL HEALTH CONCERNS PLEASE CONSULT A VET.

FOR ANY QUESTIONS REGARDING CALF XTREME, CONTACT US:

CHRIS - 027 4591 061

MATT - 021 234 1713